

MEETING ABSTRACT

Open Access

A close link between metabolic activity and functional connectivity in the resting human brain

Susanne Passow^{1,2*}, Karsten Specht^{1,3}, Tom Christian Adamsen^{4,5}, Martin Biermann^{4,6}, Njål Brekke^{4,6}, Alexander Richard Craven^{1,2}, Lars Ersland^{3,2}, Renate Grüner^{4,7,2}, Nina Kleven-Madsen^{4,7}, Ole-Heine Kvernenes⁴, Thomas Schwarzmüller^{4,8}, Rasmus Olesen⁹, Kenneth Hugdahl^{1,4,10,2}

From PSMR 2015: 4th Conference on PET/MR and SPECT/MR
La Biodola, Isola d'Elba, Italy. 17-21 May 2015

¹Department of Biological and Medical Psychology, University of Bergen, Norway

Default-mode network (DMN) functional connectivity and its task-dependent down-regulation have attracted a lot of attention in the field of neuroscience. Nevertheless, the exact underlying mechanisms of DMN functional connectivity, or more specifically, the blood oxygen level-dependent (BOLD) signal, are still not completely understood. To investigate more directly the association between local glucose consumption, local glutamatergic neurotransmission and DMN functional connectivity during rest, the present study combined for the first time 2-Deoxy-2-[18F]fluoroglucose positron emission tomography (FDG-PET), proton magnetic resonance spectroscopy (1H-MRS), and resting-state functional magnetic resonance imaging (rs-fMRI). Seed-based correlation analyses, using a key region of the DMN i.e. the dorsal posterior cingulate cortex as seed, revealed overall striking spatial similarities between fluctuations in FDG-uptake and the BOLD signal. More specifically, a conjunction analysis across both modalities showed that DMN areas as the inferior parietal lobe, angular gyrus, precuneus, middle and medial frontal gyrus were positively correlated with the dorsal posterior cingulate cortex. Furthermore, we could demonstrate that local glucose consumption in the medial frontal gyrus, posterior cingulate cortex and left angular gyrus was associated with functional connectivity within the DMN. We did not find a relationship between glutamatergic neurotransmission and functional connectivity. In line with very recent findings, our results provide further evidence for a close association between local metabolic activity and functional connectivity and enable further insights towards a better understanding of the underlying mechanisms of the BOLD signal.

Authors' details

¹Department of Biological and Medical Psychology, University of Bergen, Norway. ²NORMENT Center of Excellence, University of Oslo, Norway. ³Department of Clinical Engineering, Haukeland University Hospital, Bergen, Norway.

⁴Department of Radiology, Haukeland University Hospital, Bergen, Norway. ⁵Department of Chemistry, University of Bergen, Norway. ⁶Department of Oncology and Medical Physics, Haukeland University Hospital, Bergen, Norway.

⁷Department of Physics and Technology, University of Bergen, Norway. ⁸Department of Clinical Medicine, University of Bergen, Norway. ⁹Center of Functionally Integrative Neuroscience and MINDLab, Aarhus University, Aarhus, Denmark.

¹⁰Division of Psychiatry, Haukeland University Hospital, Bergen, Norway.

Published: 18 May 2015

doi:10.1186/2197-7364-2-S1-A78

Cite this article as: Passow et al.: A close link between metabolic activity and functional connectivity in the resting human brain. *EJNMMI Physics* 2015 **2**(Suppl 1):A78.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
